

PATIENT INFORMATION:

PATIENT'S NAME: _____ D.O.B: _____ F M
 ADDRESS: _____
 TELEPHONE #: _____ OHIP #: _____
 APPOINTMENT DATE: _____ TIME: _____ WEIGHT: _____
 *Note: Please see back for patient instructions and directions Check if applicable **STAT**

NUCLEAR CARDIOLOGY - 2300 Eglinton Ave W

MYOCARDIAL PERFUSION IMAGING - MPI
 (CARDIOLITE / MYOVUE)

Exercise Persantine (if unable to exercise)

*Note: Please consult your doctor regarding the discontinuation of Beta Blockers (48hrs), Erectile Dysfunction (48hrs), and Calcium Blockers (24hrs) prior to cardiac stressing.

AMBULATORY MONITORING

HOLTER MONITORING
 24 Hours 48 Hours
 72 Hours 2 Weeks

LOOP / CARDIAC EVENT (2 Weeks)

24HR BP MONITOR (\$80.00 fee required)

CARDIOLOGY

12-LEAD ELECTROCARDIOGRAM

ECHOCARDIOGRAM (COLOUR DOPPLER)
Please select one of the following indications:

Chest pain suspicious of CAD Murmur
 Hypertension Palpitations/arrhythmias
 Syncope Congestive heart failure
 Other: _____

EXERCISE STRESS TEST - 2300 Eglinton Ave W

CARDIOLOGY CONSULTATION

CONSULTATION REQUESTED
 Urgent, first available cardiologist
 Dr. Ali Akram
 Dr. Aman Mundi
 Dr. Abdulwahab Arrazaghi
 Dr. _____

CONSULT IF TEST RESULT IS POSITIVE / ABNORMAL

Clinical Information: _____

Referring Physician: _____ Print Name _____ Signature _____ Fax #: _____

Copy To: _____ Print Name _____ Fax #: _____

PATIENT PREPARATION AND INSTRUCTIONS

NOTES:


1. A valid health card must be shown at every visit.
2. A signed requisition by a registered physician is mandatory for all exams.
3. If you would like to cancel/ change your appointment, please notify our office at least 24 hours prior to your scheduled time.

For additional locations across Ontario, visit www.myhealthcentre.ca

NUCLEAR CARDIOLOGY

1. Patient may have a light breakfast/lunch (e.g. toast, jam, fruit, juice, water) and then nothing to eat 1 hour prior to the test.
2. Discontinue all caffeine products 24 hours prior to the test. This includes all tea, coffee, decaffeinated tea/coffee, pop, chocolate, Tylenol 2 & 3 and/or medications containing caffeine.
3. Insulin-dependent diabetics should take their insulin and a light meal 1 hour prior to the test.
4. Wear loose fitting clothing (e.g. T-shirt, track pants, athletic shoes, etc.).
5. Bring a list of all current prescription medications and check with your physician regarding the discontinuation of any heart medications (e.g. Beta-Blockers like Metoprolol or Atenolol, as well as Calcium Channel Blockers like Diltiazem or Verapamil).
6. Do not take erectile dysfunction medications (e.g. Viagra, Cialis, Levitra, etc.) 48 hours prior to the test.

MYOCARDIAL PERFUSION IMAGING consists of 2 parts: 1. Rest Study - takes approximately 1.5 - 2 hours and consists of an injection followed by imaging. 2. Stress Study - takes approximately 2 - 2.5 hours and consists of a stress test, injection and imaging.



CARDIOLOGY CLINIC

790 Burnhamthorpe Rd. W., Unit 1
 (major intersection is Burnhamthorpe and Mavis)
 Mississauga, ON L5C 2R9

Tel: 905-828-0653 | Fax: 905-878-6575